

- 11 **Lyall EGH**. Paediatric HIV in 2002—a treatable and preventable infection. *J Clin Virol* 2002;**25**:107–19.
- 12 **Sharland M**, Gibb DM, Tudor-Williams G. Advances in the prevention and treatment of paediatric HIV infection in the United Kingdom. *Arch Dis Child* 2002;**87**:178–80.
- 13 **Gibb DM**, Duong T, Tookey PA, *et al*. Decline in mortality, AIDS, and hospital admissions in perinatally HIV-1 infected children in the United Kingdom and Ireland. *BMJ* 2003;**327**:1019.
- 14 **Centers for Disease Control and Prevention**. Revised classification system for human immunodeficiency virus infection in children less than 13 years of age. *MMWR Morb Mortal Wkly Rep* 1994;**43**:1–12.
- 15 **Shann F**, Pearson G, Slater A, *et al*. Paediatric index of mortality (PIM): a mortality prediction model for children in intensive care. *Intensive Care Med* 1997;**23**:201–7.
- 16 **Sharland M**, Castelli G, Ramos JT, on behalf of the PENTA steering committee, *et al*. Penta Guidelines for the use of antiretroviral therapy in paediatric HIV infection 2001. Available at <http://www.ctu.mrc.ac.uk/penta> (accessed December 2006).
- 17 **Van Rossum AM**, Fraaij PL, de Groot R. Efficacy of highly active antiretroviral therapy in HIV-1 infected children. *Lancet Infect Dis* 2002;**2**:93–102.
- 18 **Zar HJ**, Apolles P, Argent A, *et al*. The etiology and outcome of pneumonia in human immunodeficiency virus-infected children admitted to intensive care in a developing country. *Pediatr Crit Care Med* 2001;**2**:108–12.
- 19 **Wilfret C**. Prevention of mother-to-child transmission of HIV: reflections on implementation of PMTCT in the developing world. *Acta Paediatr* 2002;**91**:863–5.
- 20 **Jeena PM**, Coovadia HM, Bhagwanjee S. Prospective controlled study of the outcome of human immunodeficiency virus-1 antibody-positive children admitted to an intensive care unit. *Crit Care Med* 1996;**24**:963–7.
- 21 **Jeena PM**, Wesley AG, Coovadia HM. Admission patterns and outcomes in a paediatric intensive care unit in South Africa over a 25-year period (1971–1995). *Intensive Care Med* 1999;**25**:88–94.
- 22 **Morris A**, Wachter RM, Luce J, *et al*. Improved survival with highly active antiretroviral therapy in HIV-infected patients with severe *Pneumocystis carinii* pneumonia. *AIDS* 2003;**17**:73–80.
- 23 **Moyle G**. Mitochondrial toxicity: myths and facts. *HIV Ther* 2004;**9**:45–47.
- 24 **Granados JMS**, Amador JTR, De Miguel SF, *et al*. Impact of highly active antiretroviral therapy on the morbidity and mortality in Spanish human immunodeficiency virus-infected children. *Pediatr Infect Dis J* 2003;**22**:863–7.
- 25 **de Martino M**, Balducci M, Galli L, *et al*. Reduction in mortality with availability of antiretroviral therapy for children with perinatal HIV-1 infection. Italian Register for HIV infection in Children and the Italian National AIDS Registry. *JAMA* 2000;**284**:190–7.
- 26 **Gortmaker SL**, Hughes M, Cervia J, *et al*. Effect of combination therapy including protease inhibitors on mortality among children and adolescents infected with HIV-1. *N Engl J Med* 2001;**345**:1522–8.
- 27 **Fauci AS**. The AIDS epidemic. Considerations for the 21st century. *N Engl J Med* 1999;**341**:1046–50.

## ARCHIVIST.....

### Hot air for head lice

Insecticide shampoos such as those that contain pyrethroids or lindane are not very effective at killing head louse eggs, may be contraindicated for patients with asthma and may give rise to resistance in head lice. Louse combs may be ineffective unless used for long periods of time. Body lice, close relatives of head lice, die when exposed for 5 min to a blow dryer delivering air at 50°C. Researchers in Salt Lake City, USA (Brad M Goates and colleagues. *Pediatrics* 2006;**118**:1962–70) have assessed six different methods of killing head lice using hot air.

The study included 169 children aged ≥6 years who had head lice and had not used a pediculocide in the past 2 weeks. The methods tested were a bonnet-style hairdryer, a handheld blow dryer with diffuse heating, a handheld blow dryer with directed heating, a wall-mounted dryer of the type used in public toilets (detached from the wall), a high volume hot air blower (the Louse-Buster) and the Louse-Buster with a hand piece combing device. The proportion of eggs killed varied from 89% with the bonnet-style hairdryer to 98% with the directed handheld blow dryer and the Louse-Buster with hand piece. The children were treated for about 30 min in their own homes, stopping temporarily if they indicated any discomfort. The proportion of lice killed varied from 10% with the bonnet-style hair dryer to 80% with the Louse-Buster with hand piece. The 11 children treated with the Louse-Buster with hand piece were followed-up after 1 week; 10 were louse-free; the other had only a single louse.

The authors of this paper recommend their custom-built machine, the Louse-Buster for use in schools but Archivist would guess that many parents might prefer to use their own hairdryers at home. Further research may be needed to determine the optimal way to use domestic hairdryers and their effectiveness.